

University of Dundee

A Morphological identification cell cytotoxicity assay using cytoplasm-localized fluorescent probe (CLFP) to distinguish living and dead cells

Lai, Fangfang; Shen, Zhengwei; Wen, Hui; Chen, Jialing; Zhang, Xiang; Lin, Ping

Published in:
Biochemical and Biophysical Research Communications

DOI:
[10.1016/j.bbrc.2016.09.169](https://doi.org/10.1016/j.bbrc.2016.09.169)

Publication date:
2017

Licence:
CC BY-NC-ND

Document Version
Peer reviewed version

[Link to publication in Discovery Research Portal](#)

Citation for published version (APA):

Lai, F., Shen, Z., Wen, H., Chen, J., Zhang, X., Lin, P., Yin, D., Cui, H., & Chen, X. (2017). A Morphological identification cell cytotoxicity assay using cytoplasm-localized fluorescent probe (CLFP) to distinguish living and dead cells. *Biochemical and Biophysical Research Communications*, 482(2), 257-263.
<https://doi.org/10.1016/j.bbrc.2016.09.169>

General rights

Copyright and moral rights for the publications made accessible in Discovery Research Portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from Discovery Research Portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain.
- You may freely distribute the URL identifying the publication in the public portal.

Take down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

A Morphological Identification Cell Cytotoxicity Assay Using Cytoplasm-localized Fluorescent Probe (CLFP) to Distinguish Living and Dead Cells

Fangfang Lai^{1,‡}, Zhengwei Shen^{2,‡}, Hui Wen³, Jialing Chen⁴, Xiang Zhang³, Ping Lin⁴, Dali Yin¹, Huaqing Cui^{1,*}, and Xiaoguang Chen^{1,*}

1, State Key Laboratory of Bioactive Substances and Function of Natural Medicine, Institute of Materia Medica, Peking Union Medical College and Chinese Academy of Medical Sciences, Beijing, 100050, China;

2, Department of Applied Mathematics, University of Science and Technology Beijing, Beijing, 100083, China;

3, Beijing Key Laboratory of Active Substances Discovery and Drugability Evaluation, Institute of Materia Medica, Peking Union Medical College and Chinese Academy of Medical Sciences, Beijing, 100050, China;

4, Department of Mathematics, University of Dundee, Dundee, DD1 4HN, UK.

Corresponding authors:

Emails: hcui@imm.ac.cn; chxg@imm.ac.cn

‡These authors contributed equally to this work.

CLFP staining

The fluorescence microscopy was performed with high content screening system (Operetta) PerkinElmer and/or fluorescence microscopy Olympus IX71.

1 Staining conditions for CLFP probe 1 and 2

HepG2 cells were cultured in 24-well plates with low density. Two CLFPs 1 and 2 were reported previously (Figure S1), the experiment condition of CLFP 1 is $25\ \mu\text{M} \times 30\ \text{min}$, UV excited; and probe 2 is $1\ \mu\text{M} \times 10\ \text{min}$, blue fluorescence excited. After incubation, the medium was washed away with PBS buffer, HepG2 cells were imaged under fluorescent microscopy. Both CLFP 1 and 2 staining can provide reliable and high resolution cell images (Figure S2)




Figure S1. Structures of two CLFPs reported in our previous study.²⁶ Probe 1 is a DNS-derived CLFP, and probe 2 is a BODIPY-derived CLFP.




Figure S2. Fluorescence microscopy images of the morphologies of HepG2 cells, which show the cytoplasm localization of probe 1 and 2. Both probe staining can give good quality cell images.

2 The imaging of fluorescent morphologies of dead cells using probe 2 staining

HepG2 cells were cultured into 24-well plates and treated with various concentrations of cytotoxicity drugs (Paclitaxel, Staurosporine) for 16 hours. The concentrations of these compounds were varied among 0.001 μM to 0.1 μM . After washed with PBS for 3 times, probe 2 was added into the wells for an hour to stain HepG2 cells, and imaged under fluorescent microscopy.

As long as the HepG2 cells were treated under the same conditions, CLFP 2 staining provides similar fluorescent images as CLFP1 staining. Figure S2 show the morphologies of dying cells as bright, solid fluorescent spheres (white arrows). The intensity of these fluorescent spheres was stronger than that of living cells. This enhanced the pan-applicability of CLFP method.




Figure S3. HepG2 cells images, which were treated with various concentrations of cytotoxicity drugs (Paclitaxel, Staurosporine). HepG2 cells then were stained with probe 2, and imaged under both fluorescence and optical microscopy. For all of the dead cells that can be observed in the optical with round shapes, we can see that they show enhanced fluorescent intensities and full shape of fluorescence in the fluorescent images. This is consistent with the CLFP 1 staining.

3. Series of paclitaxel-treated HepG2 cells were stained with CLFP 2

HepG2 cells were treated with different concentrations Paclitaxel (0, 0.0016, 0.008, 0.04, 0.2, 1 μM) for 24 hours. Cells were then stained with probe 1 for 10 min. After washing, the cells were imaged under fluorescent microscopy. Figure S4A–F show the ratio of healthy cells to dead cells in different concentrations of

paclitaxel-treated HepG2 cells. As the concentration of paclitaxel increased, fewer healthy cells were observed, whereas the number of dead cells initially increased to a maximum (Figure S4D), and then decreased (Figures S4E and S4F). This is also in agree with the CLFP 1 staining.




Figure S4. (A–F) Fluorescence microscopy images of the morphologies of HepG2 cells treated with a series of concentrations of paclitaxel.

4. Computation analysis of the fluorescent photos.




Figure S5. Anatomy of the computational analysis used to separate the areas of living and dead cells from the grayscale fluorescence microscope image.